

Although not the focus of the study, Talmud and colleagues⁹ show that even a diagnosis of a major mutation in familial hypercholesterolaemia does not always result in appropriate LDL-C treatment; the mean LDL-C concentration after treatment was only reduced by 22%, to 5.49 mmol/L (recommended goal <2.6 mmol/L). Even in the mutation-negative cohort, with lower starting LDL-C concentrations, the mean post-treatment concentration was 4.22 mmol/L, a reduction of only 28%. These results, from patients attending specialised lipid centres, are a cause for concern, especially as highly effective statins that lower LDL-C in patients with familial hypercholesterolaemia by 50% are now generic and inexpensive. Randomised trials have shown that the more LDL-C is lowered, the lower the risk for coronary artery disease. Findings from Mendelian randomisation studies¹⁰ show that polymorphisms associated with lower LDL-C starting presumably in childhood are associated with a far greater reduction in the risk of coronary heart disease than is reported in drug trials beginning later in life, after atherosclerosis has already developed.

Thus all people, irrespective of age, with raised LDL-C concentrations in whom no secondary cause can be identified, especially if they have a family history of premature coronary artery disease, should be treated for presumptive familial hypercholesterolaemia according to clinical criteria.⁵ To add the complexity of SNP analysis for minor genes and eliminate cascade LDL-C and clinical testing of relatives of patients with polygenic familial hypercholesterolaemia does not seem to be warranted, and could even be diversionary.

*Evan A Stein, Frederick J Raal

Metabolic and Atherosclerosis Research Center, Cincinnati, OH 45227, USA (EAS); and Department of Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa (FJR)
esteinmrl@aol.com

We declare that we have no conflicts of interest.

- Goldstein J, Hobbs H, Brown M. Familial hypercholesterolemia. In: Scriver C, Beauder A, Sly W, Valle D, eds. The metabolic and molecular bases of inherited disease. New York: McGraw-Hill, 2001: 2863–13.
- Usifo E, Leigh SE, Whittall RA, et al. Low-density lipoprotein receptor gene familial hypercholesterolemia variant database: update and pathological assessment. *Ann Hum Genet* 2012; **76**: 387–401.
- Slack J. Risks of ischaemic heart disease in familial hyperlipoproteinaemic states. *Lancet* 1969; **2**: 1380–82.
- Müller C. Xanthomata, hypercholesterolemia and angina pectoris. *Acta Med Scand* 1938; **89**: 75–84.
- Scientific Steering Committee on behalf of the Simon Broome Register Group. Mortality in treated heterozygous familial hypercholesterolemia: implications for clinical management. *Atherosclerosis* 1999; **42**: 105–12.
- Illingworth DR, Sexton GJ. Hypercholesterolemic effects of mevinolin in patients with heterozygous familial hypercholesterolemia. *J Clin Invest* 1984; **74**: 1972–78.
- Ellis A, Zhou R, Stein EA. Effect of lipid-lowering treatment on natural history of heterozygous familial hypercholesterolemia in past three decades. *Am J Cardiol* 2011; **108**: 223–26.
- Wierzbicki AS, Humphries SE, Minhas R. Familial hypercholesterolaemia: summary of NICE guidance. *BMJ* 2008; **337**: a1095.
- Talmud PJ, Shah S, Whittall R, et al. Use of low-density lipoprotein cholesterol gene score to distinguish patients with polygenic and monogenic familial hypercholesterolaemia: a case-control study. *Lancet* 2013; published online Feb 22. [http://dx.doi.org/10.1016/S0140-6736\(12\)62127-8](http://dx.doi.org/10.1016/S0140-6736(12)62127-8).
- Ference BA, Yoo W, Alesh I, et al. Effect of long-term exposure to lower low-density lipoprotein cholesterol beginning early in life on the risk of coronary heart disease: a mendelian randomization analysis. *J Am Coll Cardiol* 2012; **60**: 2631–39.
- Schmidt HH, Hill S, Makariou EV, Feuerstein IM, Dugi KA, Hoeg JM. Relation of cholesterol-year score to severity of calcific atherosclerosis and tissue deposition in homozygous familial hypercholesterolemia. *Am J Cardiol* 1996; **77**: 575–80.
- Voight BF, Peloso GM, Orho-Melander M, et al. Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet* 2012; **380**: 572–80.
- Ferrieres J, Lambert J, Lussier-Cacan S, Davignon J. Coronary artery disease in heterozygous familial hypercholesterolemic patients with the same LDL receptor gene mutation. *Circulation* 1995; **92**: 290–95.
- Pimstone SN, Sun XM, du Souich C, et al. Phenotypic variation in heterozygous familial hypercholesterolemia: a comparison of Chinese patients with the same or similar mutations in the LDL receptor gene in China or Canada. *Arterioscler Thromb Vasc Biol* 1998; **18**: 309–15.
- Kajinami K, Takekoshi N, Brousseau ME, Schaefer EJ. Pharmacogenetics of HMG-CoA reductase inhibitors: exploring the potential for genotype-based individualization of coronary heart disease management. *Atherosclerosis* 2004; **177**: 219–34.
- Starr B, Hadfield SG, Hutten BA, et al. Development of sensitive and specific age- and gender-specific low-density lipoprotein cholesterol cutoffs for diagnosis of first-degree relatives with familial hypercholesterolaemia in cascade testing. *Clin Chem Lab Med* 2008; **46**: 791–803.

Health in Europe—a view from across the Atlantic



From a North American perspective, it is irresistible to extend cross-national comparisons from The *Lancet* Series on Health in Europe to include the USA, particularly in light of a 2013 US National Research Council and Institute of Medicine (NRC/IOM) report¹ showing that, for most health indicators and all ages

to 75 years, the USA ranks worst or among the worst of 17 similarly affluent countries (including 15 in western Europe). The US health disadvantage was noted across diverse indicators including life expectancy (figure); prematurity and low birthweight; infant, child, and maternal mortality; incidence of and mortality from

Published Online
March 27, 2013
[http://dx.doi.org/10.1016/S0140-6736\(13\)60587-5](http://dx.doi.org/10.1016/S0140-6736(13)60587-5)
See **Series** pages 1312 and 1323




Figure: Life expectancy at birth in Europe and the USA, 2009

Source: Organisation for Economic Co-operation and Development.⁷

diabetes, cardiovascular disease, respiratory disease, infectious disease, and unintentional and intentional injury; and disability. If the USA had been included in the Series, it would have ranked close to the eastern European countries on most indicators.²

The panel that undertook the NRC/IOM study¹ faced challenges shared by the authors of this Series, including a wide array of data for many indicators from many sources, data comparability limitations, lagged health effects of upstream social policies, and difficulties in identifying discrete factors that contribute to cross-national health differences when comparing economically, politically, and culturally distinct and evolving nations. The panel concluded that there were no simple explanations and many factors were probably implicated.

An obvious difference from western Europe is the absence of universal access to medical care in the USA. The fragmentation, specialist orientation, and primary care weakness noted in the Series by Bernd Rechel and colleagues³ as likely contributors to suboptimum health in the Commonwealth of Independent States may also have contributed to some poor US outcomes. Medical care deficiencies alone, however, cannot explain the pervasive US disadvantage across such disparate indicators. Obesity prevalence undoubtedly

contributes to the high prevalence of, and mortality from, non-communicable diseases, but cannot explain high rates of HIV and sexually transmitted infections, adverse birth outcomes, gun violence, and overall injury incidence and mortality in the USA. Furthermore, studies have shown these health disadvantages even when examining only individuals of normal weight.⁴ The large ethnic and socioeconomic inequalities in health⁵ in the USA do not wholly explain the findings—disadvantages are evident when examining only white⁶ or affluent populations.^{4,7,8}

In the Series, Johan Mackenbach and colleagues⁹ conclude that the steadily increasing life expectancy in western Europe during the past four decades is a result of “combined effects of economic growth, improved health care, and successful health policies (eg, tobacco control and road traffic safety)”. Although the USA and western Europe are not hugely dissimilar in terms of overall economic growth,¹⁰ the USA has higher rates of poverty, especially child poverty, lower social mobility,¹ and greater income inequality.¹¹ Notwithstanding debates about causality,¹² income inequality has been repeatedly and strongly linked with worse population health,¹³ and, along with poverty and low social mobility, might plausibly (through varied and complex pathways) be important in the US health disadvantage. The USA has been ahead of western Europe on tobacco control^{14,15} but probably not, at least recently, on transportation safety measures, and US rates of traffic deaths are substantially higher than are those in western Europe.¹ The NRC/IOM panel recommended investigation of these potential explanations and other factors, including low participation in formal child care and preschool, weak social protections overall, and, perhaps more fundamentally as a driver of relevant policies and behaviours, a deeply rooted culture that values individualism more than social solidarity.

I hope the stark findings of the NRC/IOM report will prompt greater openness in the USA to approaches that have been associated with better results elsewhere. And as eastern and western European policy makers plan their health strategies in the face of severe economic constraints, let the shorter, sicker lives in the USA sound a cautionary note for those who would dismantle universal care systems and tear other holes in what appear, from this side of the Atlantic, to be precious social safety nets.

Paula Braveman

Center on Social Disparities in Health, University of California,
San Francisco, San Francisco, CA 94118, USA
braveman@fcm.ucsf.edu

Johan Mackenbach, one of the Series authors, and I were members of the NRC/IOM panel on understanding cross-national health differences among high-income countries. I declare that I have no conflicts of interest.

- 1 National Research Council, Institute of Medicine. US health in international perspective: shorter lives, poorer health. Washington, DC: National Academies Press, 2013.
- 2 Organisation for Economic Co-operation and Development. Health at a glance 2011: OECD indicators. Paris: OECD Publishing, 2011.
- 3 Reche B, Roberts B, Richardson E, et al. Health and health systems in the Commonwealth of Independent States. *Lancet* 2013; published online March 27. [http://dx.doi.org/10.1016/S0140-6736\(12\)62084-4](http://dx.doi.org/10.1016/S0140-6736(12)62084-4).
- 4 Crimmins EM, Preston SH, Cohen B, National Research Council. Panel on understanding divergent trends in longevity in high-income countries. International differences in mortality at older ages: dimensions and sources. Washington, DC: National Academies Press, 2010.
- 5 Braveman PA, Cubbin C, Egerter S, Williams DR, Pamuk E. Socioeconomic disparities in health in the United States: what the patterns tell us. *Am J Public Health* 2010; **100** (suppl 1): S186–96.
- 6 Banks J, Marmot M, Oldfield Z, Smith JP. Disease and disadvantage in the United States and in England. *JAMA* 2006; **295**: 2037–45.
- 7 Martinson ML, Teitler JO, Reichman NE. Health across the life span in the United States and England. *Am J Epidemiol* 2011; **173**: 858–65.
- 8 Avendano M, Glymour MM, Banks J, Mackenbach JP. Health disadvantage in US adults aged 50 to 74 years: a comparison of the health of rich and poor Americans with that of Europeans. *Am J Public Health* 2009; **99**: 540–48.
- 9 Mackenbach JP, Karanikolos M, McKee M. The unequal health of Europeans: successes and failures of policies. *Lancet* 2013; published online March 27. [http://dx.doi.org/10.1016/S0140-6736\(12\)62082-0](http://dx.doi.org/10.1016/S0140-6736(12)62082-0).
- 10 Organisation for Economic Co-operation and Development. GDP growth. National accounts at a glance 2013. Paris: OECD Publishing, 2013.
- 11 Organisation for Economic Co-operation and Development. An overview of growing income inequalities in OECD countries: main findings. Divided we stand: why inequality keeps rising. Paris: OECD Publishing, 2011.
- 12 Lynch J, Smith GD, Harper S, et al. Is income inequality a determinant of population health? Part 1. A systematic review. *Milbank Q* 2004; **82**: 5–99.
- 13 Wilkinson RG, Pickett K. The spirit level: why greater equality makes societies stronger. New York: Bloomsbury Press, 2010.
- 14 Cutler DM, Glaeser EL, National Bureau of Economic Research. What explains differences in smoking, drinking, and other health-related behaviors. Cambridge, MA: National Bureau of Economic Research, 2005.
- 15 Pampel FC. Divergent patterns of smoking across high-income nations. In: Crimmins EM, Preston SH, Cohen B, eds. International differences in mortality at older ages: dimensions and sources. Washington, DC: National Academies Press, 2010: 132–63.

Cardiology: a call for papers

Why not send your best cardiology papers to *The Lancet*? *The Lancet* is planning a special issue to coincide with the European Society of Cardiology congress to be held from Aug 31–Sept 4, 2013, in Amsterdam, The Netherlands. We will consider high quality original research papers that will influence clinical practice, and especially welcome reports that describe the results of randomised trials.

The deadline for regular submissions is May 27, 2013, via our online submission system. We will particularly welcome reports of clinical trials accepted for Hot Line sessions at the congress, for which the deadline for submissions is Aug 1.

Please state in your covering letter that the submission is in response to this call for papers. If your work is being presented at the meeting and falls under an embargo policy, please tell us the date, time, and manner of presentation (poster or oral). If your paper is accepted here, publication on *The Lancet*'s website can be scheduled to coincide with the presentation.

Stuart Spencer
The Lancet, London NW1 7BY, UK

To submit your paper go to
<http://ees.elsevier.com/thelancet>